




# Involvement of inflammasome activation via elevation of uric acid level in nociception in a mouse model of muscle pain

Molecular Pain  
Volume 15: 1–16  
© The Author(s) 2019  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1744806919858797  
journals.sagepub.com/home/mpx  


Shinichirou Yoshida<sup>1</sup> , Yoshihiro Hagiwara<sup>1</sup>, Masahiro Tsuchiya<sup>2</sup>, Masamichi Shinoda<sup>3</sup>, Masashi Koide<sup>4</sup>, Hiroyasu Hatakeyama<sup>5</sup>, Chayanit Chaweewannakorn<sup>5</sup>, Kazuaki Suzuki<sup>6</sup>, Toshihisa Yano<sup>1</sup>, Yasuhito Sogi<sup>1</sup>, Nobuyuki Itaya<sup>6</sup>, Takuya Sekiguchi<sup>7</sup>, Yutaka Yabe<sup>1</sup>, Keiichi Sasaki<sup>8</sup>, Makoto Kanzaki<sup>5</sup>  and Eiji Itoi<sup>1</sup>

## Abstract

Muscle pain is a common condition in many diseases and is induced by muscle overuse. Muscle overuse induces an increase in uric acid, which stimulates the nucleotide-binding oligomerization domain-like receptor (NLR). This receptor contains the pyrin domain NLRP-3 inflammasome which when activated, results in the secretion of potent pro-inflammatory cytokines such as interleukin-1 $\beta$  (IL-1 $\beta$ ). The aim of this study was to investigate the involvement of inflammasome activation via the elevation of uric acid level in nociception in a mouse model of muscle pain. The right hind leg muscles of BALB/c mice were stimulated electrically to induce excessive muscle contraction. The left hind leg muscles were not stimulated as a control. Mechanical withdrawal thresholds, levels of uric acid, IL-1 $\beta$ , and NLRP3, caspase-1 activity, and the number of macrophages were investigated. Furthermore, the effects of xanthine oxidase inhibitors, such as Brilliant Blue G, caspase-1 inhibitor, and clodronate liposome, on pain were investigated. In the stimulated muscles, mechanical withdrawal thresholds decreased, and the levels of uric acid, NLRP3, and IL-1 $\beta$ , caspase-1 activity, and the number of macrophages increased compared to that in the non-stimulated muscles. Administration of the inhibitors attenuated hyperalgesia caused by excessive muscle contraction. These results suggested that IL-1 $\beta$  secretion and NLRP3 inflammasome activation in macrophages produced mechanical hyperalgesia by elevating uric acid level, and xanthine oxidase inhibitors may potentially reduce over-exercised muscle pain.

## Keywords

Muscle pain, inflammasome, uric acid, electrical stimulation, mechanical hyperalgesia

Date Received: 11 September 2018; revised: 16 May 2019; accepted: 16 May 2019

## Introduction

Muscle is a common source of pain that relates to various pathologies, such as neck and shoulder pain,<sup>1</sup> non-specific lower back pain,<sup>2</sup> and myofascial pain syndrome (MPS).<sup>3</sup> Neck and shoulder pain occurs in more than 30% of the working population,<sup>4</sup> and lower back pain has a lifetime prevalence as high as 84% in the general population.<sup>5</sup> MPS is also very common in the general population, with an incidence rate as high as 50%.<sup>4</sup> Although numerous treatment methods, such as pharmacological treatments using non-steroidal

<sup>1</sup>Department of Orthopaedic Surgery, Tohoku University Graduate School of Medicine, Sendai, Japan

<sup>2</sup>Department of Nursing, Tohoku Fukushi University, Sendai, Japan

<sup>3</sup>Department of Physiology, Nihon University School of Dentistry, Tokyo, Japan

<sup>4</sup>Department of Orthopaedic Surgery, Matsuda Hospital, Sendai, Japan

<sup>5</sup>Graduate School of Biomedical Engineering, Tohoku University, Sendai, Japan

<sup>6</sup>Department of Orthopaedic Surgery, JR Sendai Hospital, Sendai, Japan

<sup>7</sup>Department of Orthopaedic Surgery, Iwate Prefectural Central Hospital, Morioka, Japan

<sup>8</sup>Division of Advanced Prosthetic Dentistry, Tohoku University Graduate School of Dentistry, Sendai, Japan

### Corresponding Author:

Yoshihiro Hagiwara, Department of Orthopaedic Surgery, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aobaku, Sendai, Japan.

Email: hagi@med.tohoku.ac.jp



anti-inflammatory drugs, physiotherapy, and dry needling, have been developed to reduce pain, no single successful strategy is available.<sup>3</sup>

Muscle overuse, especially with eccentric contraction, induces muscle pain due to an increase in adenosine triphosphate (ATP) and a lower pH.<sup>6–8</sup> Muscle fiber destruction through muscle overuse, induces the release of ATP, which activates specific receptors (so-called nociceptors and metaboreceptors).<sup>6,8</sup> Sustained muscle contraction and chronic muscle ischemia result in a decrease in pH in affected tissues.<sup>6,8</sup> Inflammatory responses to exercise-induced muscle damage also cause muscle pain.<sup>7</sup> Neutrophils and macrophages invade skeletal muscle and produce proinflammatory cytokines such as interleukin (IL)-1 $\beta$ , IL-18, and tumor necrosis factor (TNF)- $\alpha$ , after exercise-induced muscle damage.<sup>7</sup> Furthermore, we previously reported that IL-18 is secreted from neutrophils using a muscle pain model.<sup>9</sup>

Muscle overuse induces an increase in the level of uric acid, which is released from damaged cells.<sup>10–15</sup> Kono et al. reported an increase in uric acid in dead cells harvested from various tissues including the brain, heart, lung, liver, kidney, and muscle.<sup>16</sup> Uric acid is the final product of purine metabolism. In this metabolism, xanthine oxidase (XO) converts hypoxanthine to xanthine and subsequently, xanthine to uric acid.<sup>17</sup> The purine metabolism system is extremely active during high-intensity exercise and muscle ischemic conditions.<sup>11</sup> When the uric acid level in serum reaches the solubility limit, monosodium urate (MSU) crystal formation occurs,<sup>18</sup> and the crystal induces inflammasome activation.<sup>19,20</sup>

An inflammasome is a group of protein complexes that was first reported in 2002.<sup>19,21</sup> Inflammasome complexes are composed of several proteins, including the nucleotide-binding oligomerization domain-like receptor (NLR), which contains pyrin domain (NLRP) 1, NLRP3, and NLRP6. It also contains caspase activation and recruitment domain 4 (NLRC4).<sup>19,21</sup> Inflammasomes recognize inflammation-inducing stimuli and form an assembly that results in direct activation of caspase-1, which subsequently induces the secretion of potent pro-inflammatory cytokines such as IL-1 $\beta$  and IL-18, and a form of cell death called pyroptosis.<sup>19</sup> NLRP3 activates the aberrant formation of crystals from endogenous molecules such as MSU.<sup>19,20</sup> Phagocytosis of MSU crystals by macrophages and its recognition by P2 purinergic receptors result in NLRP3 inflammasome activation and release of IL-1 $\beta$ .<sup>12,19,20</sup>

IL-1 $\beta$  is a pro-inflammatory cytokine, which is released from various cells including keratinocytes, synovial cells, and macrophages.<sup>22,23</sup> Although IL-1 $\beta$  has numerous important functions under normal conditions, the overproduction of IL-1 $\beta$  is implicated in various

diseases such as rheumatoid arthritis and osteoarthritis.<sup>24,25</sup> IL-1 $\beta$  is associated with numerous states of pain.<sup>22</sup> Gout, one of the painful diseases caused by MSU deposition, is related to IL-1 $\beta$  production via NLRP3 inflammasome activation.<sup>22</sup> So et al. reported that Anakinra, a recombinant IL-1 receptor antagonist, rapidly relieved the inflammatory symptoms of gout in gouty arthritis patients.<sup>26</sup> For the onset of neuropathic pain, the interplay between the immune and nervous systems may be important, and IL-1 $\beta$  may contribute to pain.<sup>22</sup> In various animal models of neuropathic pain, IL-1 $\beta$  expression level is increased in the injured nerve, dorsal root ganglia, and spinal cord.<sup>27–29</sup> IL-1 $\beta$  is associated with persistent pain, and injury-induced central sensitization is an important mechanism for its development. IL-1 $\beta$  also acts as a mediator between glia and neurons.<sup>22,30</sup>

Involvement of inflammasome-mediated processes in numerous painful diseases including gout,<sup>20</sup> pseudogout,<sup>31</sup> and osteoarthritis<sup>32</sup> has been reported. However, few reports have described the relationship between inflammasome and muscle pain. Thus, the aim of the present study was to investigate the role of IL-1 $\beta$  secretion due to inflammasome activation in nociception in a mouse model of muscle pain.

## Materials and methods

### Experimental animals

The protocol for this experiment was approved by the Animal Research Committee of Tohoku University (approval number: 2016MdA-240). Male, 5–7-week-old BALB/c mice (body weight: 20–23 g) were obtained from Japan CLEA (Tokyo, Japan). The mice were housed under a 12:12 h light–dark cycle at 23  $\pm$  1°C.

### Repeated electrical stimulation of triceps surae muscles

Repeated electrical stimulation was used to induce excessive muscle contractions as previously described.<sup>9</sup> Briefly, two needle electrodes were applied transcutaneously into the triceps surae muscle of the right hind leg after each mouse was anesthetized with an intraperitoneal injection of medetomidine (ZENOAQ, Fukushima, Japan, 0.3 mg/kg), midazolam (SANDZ, Tokyo, Japan, 4.0 mg/kg), and butorphanol (Meiji Seika Pharma Co., Tokyo, Japan, 5.0 mg/kg). Electrical stimulation was performed at 10 Hz with a 10 V amplitude and a 100  $\mu$ s pulse width for 30 min for 7 days per week. During electrical stimulation, the right hind leg was immobilized. The ankle joint was placed in the dorsal flexion so that the triceps surae muscle was fully extended for isometric contraction. The needle electrodes were

also applied to the contralateral triceps surae muscle without electrical stimulation and immobilization.

### *Assessment of mechanical nociceptive thresholds*

Assessment of the mechanical withdrawal threshold (MWT) was performed using the Randall–Selitto test (MK-201D Pressure Analgesy-Meter, Muromachi Kikai Co., Tokyo, Japan) as previously reported.<sup>9</sup> Briefly, a linear increase in pressure (10 mm Hg/s) was applied to the lateral surface of the triceps surae muscle using a cone-shaped plastic tip attached to a scale with a display. The MWT was defined as the amount of pressure (mm Hg) required to elicit pain-related behaviors such as vocalization, struggling, and leg withdrawal.<sup>9,33,34</sup> Experiments investigating secondary hyperalgesia were performed in the plantar surface of the foot. The cut-off value of the MWT was 250 mm Hg.<sup>34</sup> On day 7 following the initiation of electrical stimulation, an assessment of the MWT data was performed in the morning as circadian rhythm affects pain sensitivity. To avoid bias, assessment of the MWT data was performed by an investigator who was blinded to the experimental conditions.

### *Local effect of MSU on hyperalgesia*

To confirm the local effects of MSU, recrystallized MSU (Monosodium Urate, No. 133–13432; Wako Pure Chemicals Industries, Osaka, Japan) dissolved in saline was administered to the right triceps surae muscle (MSU group) as previously described.<sup>35</sup> At the same time, saline (only) was administered to the contralateral triceps surae muscle (Saline group). The solution was injected under the fascia of the lateral head of triceps using a 27-gauge needle. The volume of injected solution was 50  $\mu$ l in each group. Days 0, 1, 2, and 4 post injection, MWTs were assessed and the triceps surae muscles obtained for measurements of the levels of NLRP3 and IL-1 $\beta$  by enzyme-linked immunosorbent assay (ELISA).

### *Tissue preparation*

On day 7 following the initiation of electrical stimulation, mice were sacrificed by cervical dislocation and the triceps surae muscles isolated. Specimens for ELISA and fluorometric assays were frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$ . For immunohistochemical staining, the specimens were immersed in 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.4) and stored overnight at  $4^{\circ}\text{C}$ . After dehydration through a graded series of ethanol solutions, the specimens were embedded in paraffin. The embedded tissues were then cut into 5- $\mu$ m axial sections.

### *ELISA experiments*

For the measurement of IL-1 $\beta$  levels, tissue samples were disrupted and homogenized using lysis buffer composed of bovine serum albumin (100  $\mu$ g/ml, A4503; Sigma–Aldrich, St. Louis, MO, USA), Triton X-100 (0.1%, Wako Pure Chemicals Industries), 1 M HEPES (1%, 533–08083; Wako Pure Chemicals Industries), protease inhibitor (1%, P8340; Sigma–Aldrich), and distilled water (DW). After homogenization, samples were centrifuged for 10 min at  $9,730 \times g$  and  $4^{\circ}\text{C}$ . The supernatant obtained was stored at  $-80^{\circ}\text{C}$ . IL-1 $\beta$  expression levels were analyzed with a Bio-Plex Multiplex Immunoassay System (Bio-Rad, Hercules, CA, USA) and a Bio-Plex Pro Mouse Cytokine 23-plex Assay (Bio-Rad), according to the manufacturer's instructions. Measurement of NLRP3 level was performed using Mouse NALP3/NLRP3 ELISA Kit (LS-F17336, LifeSpan Biosciences Inc., Seattle, WA, USA). Tissue samples were disrupted and homogenized using phosphate-buffered saline (PBS). Following homogenization, the samples were repeatedly frozen ( $-20^{\circ}\text{C}$ ) and thawed three times for cell lysing at room temperature. The samples were then centrifuged for 5 min at  $5,000 \times g$  and  $4^{\circ}\text{C}$ . The supernatant was utilized for the assay, according to the manufacturer's instructions.

### *Fluorometric assay experiments*

Measurement of the level of uric acid and caspase-1 activity was performed using Uric Acid Colorimetric/Fluorometric Assay Kit (K608-100, BioVision Inc., Milpitas, CA, USA) and Caspase 1 Assay Kit (ab39412, Abcam plc), respectively. Tissue samples were disrupted and homogenized using buffers from each kit. After homogenization, the samples were centrifuged for 5 min at 12,000 rpm and  $4^{\circ}\text{C}$ . The supernatant was used for the assay according to the manufacturer's instructions. Values of caspase-1 activity were normalized to the controls which were not stimulated and administered any drugs.

### *Immunohistochemistry*

Tissue sections were deparaffinized and washed in PBS. They were subsequently incubated with a solution containing Proteinase K (Takara Bio Inc. Shiga, Japan, 25  $\mu$ l), 0.5 M ethylenediaminetetraacetic acid (Invitrogen, Carlsbad, CA, USA, 0.5 ml), 1 M Tris-Cl (pH 8.0, 2.5 ml) and 50 ml DW, for 5 minutes at  $37^{\circ}\text{C}$  to induce antigen retrieval. After washing in PBS, endogenous immunoglobulins were blocked by incubation with 10% normal goat serum (Nichirei Biosciences Inc., Tokyo, Japan) for 3 h. The slides were once again washed with PBS and incubated with a polyclonal rabbit anti-mouse NLRP3 antibody (NBP2-12446,

Novus Biologicals, Littleton, CO, USA; dilution, 1:25), polyclonal rabbit anti-mouse caspase-1 antibody (ab1872, Abcam plc, Cambridge, UK, dilution 1: 25), polyclonal rabbit anti-mouse IL-1 $\beta$  antibody (ab9722, Abcam plc, concentration of 10  $\mu$ g/ml) and a monoclonal rat anti-mouse Cluster of Differentiation (CD) 68 antibody (ab53444, Abcam plc, concentration of 10  $\mu$ g/ml) in PBS overnight at 4°C. PBS was then used to rinse the slides. Subsequently, the slides were incubated for 1 h in PBS with an Alexa Fluor 488-conjugated goat anti-rabbit IgG (A-11034, Life Technologies, Carlsbad, CA, USA; dilution, 1:750) for NLRP3, caspase-1, IL-1 $\beta$  and an Alexa Fluor 555-conjugated goat anti-rat IgG (A-21434, Life Technologies; dilution, 1:750) for CD68 at room temperature. The slides were once again rinsed with PBS. Finally, the slides were incubated with 4,6-diamidino-2-phenylindole (Sigma–Aldrich; dilution, 1:500) for 10 min at 25°C for nuclear staining. Images were captured using a fluorescence microscope (BZ-9000 Bioevo, Keyence, Osaka, Japan). The images were analyzed using Adobe Photoshop (Adobe System Inc., San Jose, CA, USA). At least three images in each slide were captured at 200  $\times$  magnification and the number of CD68-positive cells (macrophages) counted. The number was presented as cells/view. To avoid bias, the evaluation was performed by two investigators who were blinded to the experimental conditions. Two animals were used for immunohistochemistry, and two slides/animal were analyzed. After confirming reproducibility, representative images were presented.

### Assessment of the effect of drug administration

Several agents were administered intraperitoneally during repeated electrical stimulations of the triceps surae muscles. This was to confirm the suppressing effects of hyperalgesia. Allopurinol (A8003, Sigma–Aldrich; 200 mg/kg/72 h),<sup>36</sup> Febuxostat (F0847, Tokyo Chemical Industry Co., Ltd., Tokyo, Japan; 5 mg/kg/24 h),<sup>37</sup> Brilliant Blue G pure (BBG, B0770, Sigma–Aldrich; 45.5 mg/48 h),<sup>38</sup> Caspase-1 inhibitor Z-Trp-Glu(OMe)-His-Asp(OMe)-fluoromethylketone (Z-WEHD-FMK, FMK002, R&D Systems, Inc., Minneapolis, MN, USA; 1 mg/kg/24 h),<sup>39</sup> and Liposomal clodronate (Xyglia Bioscience, Osaka, Japan; 200  $\mu$ l/body/48 h)<sup>40</sup> were used.

Allopurinol and Febuxostat are XO inhibitors which reduce uric acid formation by inhibiting XO which converts hypoxanthine to xanthine and uric acid.<sup>36,37</sup> BBG is a selective antagonist that attenuates NLRP3 inflammasome activation.<sup>38</sup> Caspase-1 inhibitor Z-WEHD-FMK is a caspase-1 inhibitor used to block caspase-1 activity and subsequently, the production of IL-1 $\beta$ .<sup>39</sup> Clodronate liposome induces macrophage depletion by killing these cells as a result of accumulation and irreversible metabolic damage.<sup>40</sup> Control animals received an equivalent volume of 0.9% saline (saline group).

The MWTs were assessed by performing the Randall–Selitto test on day 7. Measurement of uric acid, NLRP3, IL-1 $\beta$  levels and caspase activity was performed as described above.

### Assessment of hyperalgesia in IL-1 knock out mouse

To confirm IL-1 $\beta$  effects on hyperalgesia, homozygous IL-1 knock out (KO) mice were used.<sup>41</sup> Repeated electrical stimulation was performed on IL-1 KO mice as described above, and the MWTs were assessed by performing the Randall–Selitto test on day 7.

### Statistical analysis

Statistical analysis was performed using SPSS Statistics 24 (IBM, Armonk, NY, USA). Analysis of the MWT time-course data was performed using 2-way analysis of variance (ANOVA), and repeated measurements compared by the Tukey's post-hoc multiple-comparison test. To compare data from more than three groups from a single day (MWTs, ELISA, and fluorometric assay), 1-way ANOVA with Tukey's post-hoc multiple-comparison test was used for the analysis. The NLRP3 and IL-1 $\beta$  data between the two groups were analyzed using the Wilcoxon signed-rank test. Other data between two groups were analyzed through the use of the paired t test. All data are expressed as the mean  $\pm$  standard error of the mean. A p value < 0.05 was considered statistically significant.

## Results


### MSU induced mechanical hyperalgesia and inflammasome activation

The MWTs of the MSU group significantly decreased from day 1 following the administration, compared to that before administration. Decrease in MWTs continued at least 4 days after administration. In the saline group, no significant changes were observed during the experimental periods compared to that before administration. Significant differences were observed between the two groups between 1 to 4 days after administration (Figure 1(a)). On days 2 and 4 following administration, NLRP3 and IL-1 $\beta$  induced in the MSU group significantly increased compared to the saline group based on ELISA data (Figure 1(b) and (c)).

### Excessive contraction induced mechanical hyperalgesia, elevated uric acid level, and activated inflammasome in muscles

The MWTs of the stimulated muscles significantly decreased compared to that of non-stimulated contralateral muscle (Figure 2(a)). There was no evidence for





**Figure 1.** Change in mechanical withdrawal thresholds (MWTs), NLRP3 and IL-1 $\beta$  levels by local administration of MSU. Time course of MWT values in the MSU group and saline group (a). The MWTs of the MSU group significantly decreased compared to those of the saline group, 1 to 4 days after administration. The quantities of NLRP3 (b) and IL-1 $\beta$  (c) in muscle tissues. The protein expression, determined by ELISA, increased significantly in MSU groups compared to saline groups. \* $p < 0.05$ : significantly different from the saline group; # $p < 0.05$ : significantly different from the preinjection values; The protein concentration for all samples used in the ELISA was 1.1 mg/ml. MSU: monosodium urate. IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.

secondary hyperalgesia during the experiments. Induction of uric acid, NLRP3, caspase-1 activity, and IL-1 $\beta$  in the stimulated muscles significantly increased compared to that in non-stimulated muscles (Figure 2 (b) to (e)).

#### *Inflammasome activation in macrophages in stimulated muscles*


Immunohistochemical staining revealed an increased number of CD68-positive cells (macrophages) in the stimulated muscles compared to those that were not stimulated. Most of these cells were positive for NLRP3, caspase-1, and IL-1 $\beta$  (Figure 3(a) to (r)). The number of cells that were positive for both IL-1 $\beta$  and CD68 in the stimulated muscles significantly increased compared to that of non-stimulated muscles (Figure 3(s)). In the clodronate liposome group, the MWTs significantly increased while NLRP3, caspase-1 activity, and IL-1 $\beta$  levels significantly decreased compared to those in the saline group (Figure 3(t) to (w)). No significant differences were observed between the saline group and naïve group with electrical stimulation.

#### *XO inhibitors (allopurinol and febuxostat) attenuated hyperalgesia in excessively contracted muscles*

In both the allopurinol and febuxostat groups, the MWTs significantly increased while uric acid, NLRP3, caspase-1 activity, and IL-1 $\beta$  levels significantly decreased compared to those in the saline group (Figure 4(a) to (e)). No significant differences were observed between the saline group and the naïve group with electrical stimulation. The levels of uric acid and caspase-1 activity of the allopurinol and febuxostat group did not decrease to the same level as the group without stimulation.

#### *BBG attenuated hyperalgesia in excessively contracted muscles*

The MWTs significantly increased while NLRP3, caspase-1 activity, and IL-1 $\beta$  levels significantly decreased in the BBG group compared to that in the saline group (Figure 5(a) to (d)). No significant differences were observed between the saline and the naïve groups with electrical stimulation. The caspase-1 activity of the BBG group did not decrease to the same level as the group without stimulation.



**Figure 2.** Changes in the MWTs and levels of uric acid, NLRP3, caspase-1, and IL-1 $\beta$  in stimulated and nonstimulated muscles at seven days after initiating electrical stimulation. The MWTs of stimulated muscles significantly decreased compared to those of nonstimulated contralateral muscles (a). The quantities of uric acid (b), NLRP3 (c), IL-1 $\beta$  (e), and fluorescence intensity of caspase-1 activity (d) in muscle tissues. These levels increased significantly in stimulated muscles compared to nonstimulated muscles; the concentrations of NLRP3 and IL-1 $\beta$  were equalized to 1.1 mg/ml. The concentration of uric acid is shown as nmol/50  $\mu$ l and the fluorescence intensity of caspase-1 activity shown as a ratio to the value of nonstimulated muscles. IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.

### Z-WEHD-FMK attenuated hyperalgesia in excessively contracted muscles

In the Z-WEHD-FMK group, the MWTs significantly increased while caspase-1 activity and IL-1 $\beta$  levels significantly decreased compared to the saline group (Figure 6(a) to (c)). No significant differences were observed between the saline and the naïve groups with electrical stimulation. The caspase-1 activity of the Z-WEHD-FMK group did not decrease to the same level as the group without stimulation.

### Excessive contraction did not induce mechanical hyperalgesia in IL-1 KO mice

The MWTs of the stimulated muscles did not decrease compared to those in the non-stimulated contralateral muscle in the IL-1 KO mice (Figure 7(a)). In the IL-1 KO mice, NLRP3 level of the stimulated muscle significantly increased compared to the non-stimulated muscle. Although there was no significant difference, the level of caspase-1 activity in the stimulated muscle tended to increase. For IL-1 KO mice, these levels were lower than those of naïve mice (Figure 7(b) to (c)).

## Discussion

The present study revealed the following: 1. The administration of MSU in the triceps surae muscles induced hyperalgesia and IL-1 $\beta$  elevation owing to

inflammasome activation, 2. Excessive muscle contraction by electrical stimulation induced mechanical hyperalgesia, elevated uric acid, and activated inflammasome and IL-1 $\beta$  in the muscle, 3. The number of macrophages increased after electrical stimulation and NLRP3 inflammasome activation in these macrophages, 4. Intraperitoneal administration of drugs including XO inhibitors, BBG and Z-WEHD-FMK attenuated hyperalgesia caused by excessive muscle contraction, similar to IL-1 KO mice. The pathway of muscle pain in our model and the effects of inhibitors at each stage are presented in Figure 8.


In our previous study, repeated muscle contraction (7 days) produced mechanical hyperalgesia on days 4–11 following the initiation of contraction; recovery occurred after day 12.<sup>9</sup> In the present study, which was conducted using the same model, MWTs of the stimulated muscles significantly decreased compared to those of non-stimulated contralateral muscles on day 7. Therefore, the stimulated muscles which were harvested on day 7 was appropriate for investigation of muscle pain induced by repeated excessive contraction. Although needle electrodes were also inserted into the contralateral muscle without electrical stimulation, hyperalgesia of the muscle was not observed. This finding demonstrates that the observed muscle hyperalgesia was caused by electrical stimulation, and not by the mechanical irritation of muscle tissue caused by insertion of the needle. In the present study, the muscle on the

contralateral side was used as a control (internal control) to exclude individual differences in mice. If the muscle on the contralateral side is compared to the same muscle in another mouse (external control), some differences should affect the results because the number of circulating leukocytes increases after exercise,<sup>7</sup> which must be dependent on individual specificity.

MSU activates NLRP3 inflammasome.<sup>12,19,20</sup> There are several reports that evaluate inflammasome activation by local administration of MSU.<sup>35,42,43</sup> Ju et al. and Yang et al. reported MSU injection into the soles of mice

hindlimb feet, induced elevation of caspase-1 activity and IL-1 $\beta$ .<sup>35,42</sup> Lee et al. reported elevation of NLRP3, caspase-1 activity and IL-1 $\beta$  by MSU injection into the subcutaneous tissue in the back of mice.<sup>43</sup> The data in this study was seen to correspond with these studies.

Muscle overuse induces an increase in uric acid level in human and mouse.<sup>10,11,13-15</sup> Balsom et al. and Jówko et al. reported an increase in uric acid level in serum after sprint in physical education students.<sup>10,11</sup> Chatzinikolaou et al. and Andersson et al. reported that there is an elevated blood uric acid concentration



**Figure 3.** The number of macrophages in muscle, with or without electrical stimulation, and the effect of clodronate liposome (CL) administration. Immunohistochemical staining of NLRP3 (green), caspase-1 (green), IL-1 $\beta$  (green), and CD68 (red) in stimulated and nonstimulated muscles on day 7 is shown ((a) to (r)). The number of CD68-positive cells (macrophages) in stimulated muscles increased and most of these cells are also positive for NLRP3, caspase-1, and IL-1 $\beta$ ; scale bar = 50  $\mu$ m. The number of cells copositive for IL-1 $\beta$  and CD68 increased significantly in stimulated muscles (s). The MWTs increased (t), and levels of NLRP3 (u), caspase-1 activity (v), and IL-1 $\beta$  (w) significantly decreased in the CL group compared to those in the stimulated muscle of the saline group and naïve group. The concentrations of NLRP3 and IL-1 $\beta$  are equalized to 1.1 mg/ml. The fluorescence intensity of caspase-1 activity is shown as a ratio with the value of nonstimulated muscles in naïve group. IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.





Figure 3. Continued.


and muscle soreness after exercise in team handball players and elite female soccer players, respectively.<sup>13,15</sup> Retamoso et al. reported an increase in uric acid level in the gastrocnemius muscle of mouse after downhill running exercise.<sup>14</sup> These reports supported our findings

that an increase in uric acid in the stimulated muscles presented hyperalgesia. Although an increase in uric acid level and NLRP3 inflammasome activation was observed in stimulated muscles, MSU crystal formation was not confirmed as the polarizing microscope could






**Figure 4.** Changes in MWTs and the levels of uric acid, NLRP3, caspase-1 activity, and IL-1 $\beta$  produced after administration of xanthine oxidase (XO) inhibitors. The MWTs increased (a), and levels of uric acid (b), NLRP3 (c), caspase-1 activity (d), and IL-1 $\beta$  (e) significantly decreased in the allopurinol group and febuxostat group when compared to the stimulated muscles of the saline group and the naïve group. The concentrations of NLRP3 and IL-1 $\beta$  are equalized to 1.1 mg/ml and the concentration of uric acid presented as nmol/50  $\mu$ l. The fluorescence intensity of caspase-1 activity is shown as a ratio with the value of nonstimulated muscles in the naïve group. IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.



**Figure 5.** Changes in MWTs and the levels of NLRP3, caspase-1 activity, and IL-1 $\beta$  produced after administration of brilliant blue G (BBG). The MWTs increased (a), and levels of NLRP3 (b), caspase-1 activity (c), and IL-1 $\beta$  (d) significantly decreased in the BBG group compared to the stimulated muscles of the saline group and the naïve group. The concentrations of NLRP3 and IL-1 $\beta$  are equalized to 1.1 mg/ml. The fluorescence intensity of caspase-1 activity is shown as a ratio to the value of nonstimulated muscles in the naïve group. IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.




**Figure 6.** Changes in MWTs and the levels of caspase-1 activity and IL-1 $\beta$  produced after administration of Z-Trp-Glu(OMe)-His-Asp(OMe)-fluoromethylketone (Z-WEHD-FMK). The MWTs increased (a), and levels of caspase-1 activity (b) and IL-1 $\beta$  (c) significantly decreased in the Z-WEHD-FMK group compared to the stimulated muscle of saline group and naive group. The concentrations of IL-1 $\beta$  were equalized to 1.1 mg/ml. The fluorescence intensity of caspase-1 activity is shown as a ratio with the value of nonstimulated muscles in the naive group. IL: interleukin.

not be used. According to previous studies, the solubility limit of uric acid is 405  $\mu\text{mol/l}$  in serum.<sup>18</sup> In our results, the average of local uric acid concentration in stimulated muscle was almost 600  $\mu\text{mol/l}$ , which is sufficient to enable crystal formation. Therefore, NLRP3 inflammasome activation in the stimulated muscles must have occurred.


Although there are very few reports of the direct relationship between NLRP3 inflammasome and muscle pain, several studies have reported the relationship between NLRP3 inflammasome and fibromyalgia, and chronic fatigue syndrome, which result in muscle pain and fatigue.<sup>44–46</sup> Bullón et al. and Cordero et al. reported that overactivation of NLRP3 inflammasome occurs in blood mononuclear cells in patients with fibromyalgia.<sup>44,45</sup> Zhang et al. reported the activation of NLRP3 inflammasome in the diencephalons, which is responsible for fatigue sensation. This experiment was conducted using a mouse fatigue model with lipopolysaccharide (LPS) combined with swim stress.<sup>46</sup> These previous reports may support our findings of the increase in NLRP3, caspase-1 activity and IL-1 $\beta$  in the stimulated muscles. Although evaluation of NLRP3 inflammasome activity was performed only in peripheral muscle tissues without any testing of the central nervous system (CNS) in this study, it is quite possible that

NLRP3 inflammasome activation also occurs in the CNS. This is based on the study by Zhang et al.<sup>46</sup>

There are a few reports of NLRP3 inflammasome activation in skeletal muscle cell.<sup>47,48</sup> Ding et al. reported NLRP3 inflammasome activation in interferon gamma (IFN- $\gamma$ ) treated C2C12 cultured myotubes.<sup>47</sup> McBride et al. reported that NLRP3 inflammasome contributes to sarcopenia.<sup>48</sup> However, to the best of our knowledge, there is no previous report of the involvement of NLRP3 inflammasome in muscle pain. NLRP3 is mainly expressed by myeloid cells including monocytes and macrophages.<sup>12</sup> In addition, the activation of NLRP3 inflammasome induced by uric acid occurs in macrophages.<sup>19,20</sup> Martinon et al. and Gicquel et al. reported inflammasome activation in monocyte cell line THP1 and human macrophages which were incubated with MSU crystals.<sup>12,20</sup> These reports support our findings that NLRP3 inflammasome activation occurred in macrophages which invaded the stimulated muscle. There are a few reports that clodronate liposome depletes macrophages and suppresses NLRP3 inflammasome activation and IL-1 $\beta$  release in several disease models such as lung tumor and alcoholic hepatosteatosis.<sup>49,50</sup> As previously mentioned, no report has addressed muscle pain. Gregory et al. reported that clodronate liposome reduced hyperalgesia induced by injection of normal



**Figure 7.** Changes in MWTs and the levels of NLRP3 and caspase-1 activity in naïve mice and IL-1 knock-out mice. A significant difference was not observed in the MWTs between stimulated and nonstimulated muscles in IL-1 KO mice (a). In the IL-1 KO mice, the levels of NLRP3 (b) and caspase-1 activity (c) in the stimulated muscle tended to increase compared to levels in the nonstimulated muscle. The concentrations of NLRP3 are equalized to 1.1 mg/ml. The fluorescence intensity of caspase-1 activity is shown as a ratio to the value of nonstimulated muscles in the naïve group. KO: knock out; IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain.



**Figure 8.** Schema of the pathway of muscle pain in the mouse model, and the effects of inhibitors at each stage. MSU: monosodium urate; KO: knock out; IL: interleukin; NLRP: nucleotide-binding oligomerization domain-like receptor pyrin domain; Z-WEHD-FMK: Z-Trp-Glu (OMe)-His-Asp(OMe)-fluoromethylketone.

saline adjusted to pH 5.0 into muscle when combined with an electrical stimulation. They concluded that muscle fatigue decreases the pH of muscle and activates acid-sensing ion channel 3 (ASIC3) in macrophages.

This in turn enhances hyperalgesia of the muscle.<sup>51</sup> Although the detailed mechanisms are different, their report suggests that macrophages are involved in muscle pain and this corresponds with our results.

There are several previous reports that XO inhibitors inhibit NLRP3 inflammasome activation in various disease models.<sup>52,53</sup> Aibibula et al. reported that febuxostat inhibited NLRP3 inflammasome activation in the mouse model of metabolic syndrome.<sup>52</sup> Similarly, Wan et al. reported that allopurinol inhibited NLRP3 inflammasome activation in the mouse model of non-alcoholic fatty liver disease.<sup>53</sup> Thus, these reports supported our findings that allopurinol and febuxostat inhibited NLRP3 inflammasome activation in the stimulated muscle. In the present study, although the uric acid levels in the allopurinol group and febuxostat group did not decrease to the level seen in non-stimulated muscles, the MWT was recovered to the same level as that of the non-stimulated muscle. There are a few studies suggesting that different mechanisms for uric acid-mediated NLRP3 inflammasome activation is the cause of pain.<sup>36,54</sup> Schmidt et al. reported that administration of allopurinol produced dose-dependent antinociceptive effects in chemical and thermal pain models.<sup>36</sup> They concluded that allopurinol-induced anti-nociception may be related to adenosine accumulation which induces a decrease in the release of painful substances including substance P and glutamate.<sup>36</sup> Ives et al. reported that XO-derived reactive oxygen species, excluding uric acid, is the trigger for NLRP3 inflammasome activation and XO blockade impair it.<sup>54</sup> These mechanisms may explain the dissociation observed in the present study. Allopurinol and febuxostat are used clinically to treat hyperuricemia and gout.<sup>36,37</sup> Since allopurinol and its active metabolite are excreted by the kidneys, a life-threatening toxicity syndrome consisting of an erythematous, desquamative skin rash, fever, hepatitis, eosinophilia, and worsening in renal function can occur if used by patients with renal insufficiency.<sup>55,56</sup> In contrast, febuxostat can be more safely administered to such patients because it does not affect renal excretion.<sup>56</sup> As the safety of these drugs needs to be confirmed before use, it is helpful to use existing medication for chronic pain.

BBG is a selective P2X7 receptor antagonist that attenuates NLRP3 inflammasome activation.<sup>38</sup> The P2X7 receptor is a trimeric ion channel gated by extracellular ATP which present in numerous types of cell including stem, blood, glial, neural, bone, endothelial, muscle, renal and skin cells.<sup>57</sup> BBG attenuates various types of pain including muscle pain and neuropathic pain.<sup>9,58</sup> In our previous study, BBG attenuated hyperalgesia caused by excessive muscle contraction.<sup>9</sup> There are several reports that BBG inhibits NLRP3 inflammasome activation in various disease models.<sup>38,59</sup> Wang et al. reported BBG inhibited NLRP3 inflammasome activation in the bone marrow derived from macrophages. This was observed using an LPS-induced acute lung injury mouse model.<sup>38</sup> Furthermore, Zhong et al.

reported inhibition of NLRP3 inflammasome activation by BBG using graft-versus host-disease model mice produced by allogeneic hematopoietic stem cell transplantation.<sup>59</sup> These reports support our findings that BBG inhibited NLRP3 inflammasome activation in the stimulated muscle. BBG has already been used in clinical settings such as in ophthalmic surgery and confirmed a safe and reliable dye.<sup>60</sup> However, there is no previous report investigating the effects of systemic administration of BBG on the human body.

Z-WEHD-FMK is a caspase-1 inhibitor that inhibits the caspase-1 activity and the subsequent production of IL-1 $\beta$ .<sup>39</sup> There are a few reports that caspase-1 inhibitor attenuated pain.<sup>23,61</sup> Li et al. reported caspase-1 inhibitor attenuated mechanical allodynia in the rat model of complex regional pain syndrome type I.<sup>23</sup> Chen et al. reported caspase-1 inhibitor reduced IL-1 $\beta$  level in a rat model of headache induced by intrathecal injection of inflammatory chemicals.<sup>61</sup> These reports support our findings that Z-WEHD-FMK attenuated pain and reduced IL-1 $\beta$  level in the stimulated muscle. Although several reports have investigated the effect of Z-WEHD-FMK on human cells *in vitro*,<sup>62,63</sup> there is no previous report investigating the effects of Z-WEHD-FMK administration to humans.

In IL-1 KO mice, although NLRP3 and caspase-1 activity levels in the stimulated muscle increased compared to the non-stimulated muscle, the MWTs of these groups were almost equal. This indicates that IL-1 $\beta$ , and not NLRP3 or caspase-1, is involved in the development of muscle pain. In the IL-1 KO mice, NLRP3 and caspase-1 activity levels were lower than in naïve mice. Zaslona et al. reported that pro-IL-1 $\beta$  production induces mature IL-1 $\beta$  production via NLRP3 activation in macrophages (i.e., there is a feed-forward loop among IL-1 $\beta$  production via NLRP3 activation).<sup>64</sup> This may explain the lower levels of NLRP3 and caspase-1 activity in IL-1 KO mice compared to those in naïve mice.

From the results of the drug administration tests, caspase-1 activities of treated muscle increased when compared to those of non-stimulated muscle. The activation of inflammasomes occur via various proteins such as NLRP1 and NLRC4, in addition to NLRP3.<sup>19</sup> Recognition of signals by these inflammasome proteins results in the activation of caspase-1.<sup>19</sup> Therefore, caspase-1 activation through other inflammasome proteins than NLRP 3 may occur and might have been the cause of electrode puncture or electrical stimulation.

In the present study, IL-1 KO mice, which are deficient in IL-1 $\alpha$  and IL-1 $\beta$  genes, were used instead of the IL-1 $\beta$  KO mice to confirm the effect of IL-1 $\beta$  on hyperalgesia. This is because the activation of NLRP3 inflammasome can induce the secretion of not only IL-1 $\beta$ , but also IL-1 $\alpha$ .<sup>65,66</sup> If only the IL-1 $\beta$  gene was deficient, the level of IL-1 $\alpha$  can increase as such cytokines display



mutual compensation.<sup>67</sup> The physiological role of IL-1 $\alpha$  is not well-defined<sup>65</sup>; however, several studies reported that the progression of inflammatory diseases may not solely be due to IL-1 $\beta$ , but also IL-1 $\alpha$ .<sup>66,68</sup> Therefore, IL-1 KO mice were used in this study.

Numerous drugs were used to inhibit inflammasome activation at various stages in the present study. XO inhibitors reduced uric acid formation, P2X7 receptor antagonist attenuated NLRP3 inflammasome activation, and caspase-1 inhibitor attenuated caspase-1 activity. Ultimately, these drugs attenuated the production of IL-1 $\beta$  and pain. Therefore, these drugs may potentially reduce over-exercised muscle pain in humans. In particular, the XO inhibitors can be used to treat human muscle pain as they are already safely used in the human body.

Sjøgaard et al. reported that there are differences between muscle pain such as shoulder and neck pain, which occurs during occupational tasks, and pain that occurs during physical activities at leisure and sports.<sup>69</sup> The former is induced by static sustained and monotonous repetitive muscle contractions, despite a rather low relative muscle load.<sup>69</sup> In contrast, the latter is induced by more dynamic and relatively high muscle forces.<sup>69</sup> As the possibility remains that our model could not completely produce the pathology of chronic muscle pain such as shoulder and neck pain, lower back pain, and MPS, further research is necessary to investigate appropriate electrical stimulation conditions to reflect the pathology of chronic muscle pain. Other limitations are as follows: (1) the evaluation of inflammasome activation and IL-1 $\beta$  levels was performed only in muscle tissues; no additional testing of sensory neurons and the CNS was performed, (2) the evaluation of tissue pH and ATP production was not performed, (3) we did not confirm MSU crystal formation using polarizing microscopes, (4) the cut-off value of uric acid concentration that can induce inflammasome activation was not determined, (5) an analysis of other inflammatory cytokines such as IL-6, TNF- $\alpha$ , and IFN- $\gamma$ , was not performed, and (6) the evaluation of inflammasome activation was not performed with neutrophils.

## Conclusions

IL-1 $\beta$  secretion and NLRP3 inflammasome activation in macrophages due to elevated levels of uric acid produced mechanical hyperalgesia through repeated excessive muscle contractions. Therefore, pharmacological blockade of this process may potentially reduce over-exercised muscle pain in humans.

## Acknowledgments

The authors thank Shunji Sugawara (Division of Oral Immunology, Tohoku University Graduate School of Dentistry, Sendai, Japan) for providing IL-1 KO mice and

Natsumi Emoto (Graduate School of Biomedical Engineering, Tohoku University, Sendai, Japan) for technical advice and data collection.

## Author Contributions

Shinichirou Yoshida performed experiments and wrote the article. Yoshihiro Hagiwara and Masahiro Tsuchiya supervised the research design and writing of the article. Masamichi Shinoda and Makoto Kanzaki supervised the research design and collection of data. Masashi Koide, Hiroyasu Hatakeyama, Kazuaki Suzuki, and Chayanit Chaweewannakorn participated in data collection. Toshihisa Yano, Yasuhito Sogi, Nobuyuki Itaya, Takuya Sekiguchi, and Yutaka Yabe gave advice regarding the experimental methods used. Keiichi Sasaki and Eiji Itoi provided advice regarding research design.



## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## ORCID iD

Shinichirou Yoshida  <https://orcid.org/0000-0003-1110-2363>  
Makoto Kanzaki  <https://orcid.org/0000-0002-6884-2955>

## References

1. Hanvold TN, Wærsted M, Mengshoel AM, Bjertness E, Stigum H, Twisk J, Veiersted KB. The effect of work-related sustained trapezius muscle activity on the development of neck and shoulder pain among young adults. *Scand J Work Environ Health* 2013; 39: 390–400.
2. van Dieen JH, Selen LP, Cholewicki J. Trunk muscle activation in low-back pain patients, an analysis of the literature. *J Electromyogr Kinesiol* 2003; 13: 333–351.
3. Borg-Stein J, Simons DG. Focused review: myofascial pain. *Arch Phys Med Rehabil* 2002; 83: S40–S47, s48–s49.
4. Punnett L, Wegman DH. Work-related musculoskeletal disorders: the epidemiologic evidence and the debate. *J Electromyogr Kinesiol* 2004; 14: 13–23.
5. Balague F, Mannion AF, Pellise F, Cedraschi C. Non-specific low back pain. *Lancet (London, England)* 2012; 379: 482–491.
6. Mense S. Muscle pain: mechanisms and clinical significance. *Dtsch Arztebl Int* 2008; 105: 214–219.
7. Peake J, Nosaka K, Suzuki K. Characterization of inflammatory responses to eccentric exercise in humans. *Exerc Immunol Rev* 2005; 11: 64–85.
8. Mastaglia FL. The relationship between muscle pain and fatigue. *Neuromuscul Disord* 2012; 22: S178–S180.
9. Yoshida S, Hagiwara Y, Tsuchiya M, Shinoda M, Koide M, Hatakeyama H, Chaweewannakorn C, Yano T, Sogi

- Y, Itaya N, Sekiguchi T, Yabe Y, Sasaki K, Kanzaki M, Itoi E. Involvement of neutrophils and interleukin-18 in nociception in a mouse model of muscle pain. *Mol Pain* 2018; 14:1744806918757286.
10. Balsom PD, Seger JY, Sjodin B, Ekblom B. Physiological responses to maximal intensity intermittent exercise. *Europ J Appl Physiol* 1992; 65: 144–149.
  11. Jowko E, Dlugolecka B, Makaruk B, Cieslinski I. The effect of green tea extract supplementation on exercise-induced oxidative stress parameters in male sprinters. *Eur J Nutr* 2015; 54: 783–791.
  12. Gicquel T, Robert S, Loyer P, Victoni T, Bodin A, Ribault C, Gleonnet F, Couillin I, Boichot E, Lagente V. IL-1 $\beta$  production is dependent on the activation of purinergic receptors and NLRP3 pathway in human macrophages. *Faseb J* 2015; 29: 4162–4173.
  13. Chatzinikolaou A, Christoforidis C, Avloniti A, Draganidis D, Jamurtas AZ, Stampoulis T, Ermidis G, Sovatzidis A, Papassotiropoulos I, Kambas A, Fatouros IG. A microcycle of inflammation following a team handball game. *J Strength Cond Res* 2014; 28: 1981–1994.
  14. Retamoso LT, Silveira MEJ, Lima FD, Busanello GL, Bresciani G, Ribeiro LR, Chagas PM, Nogueira CW, Braga AC, Furian AF, Oliveira MS, Figuera MR, Royes LF. Increased xanthine oxidase-related ROS production and TRPV1 synthesis preceding DOMS post-eccentric exercise in rats. *Life Sci* 2016; 152: 52–59.
  15. Andersson H, Raastad T, Nilsson J, Paulsen G, Garthe I, Kadi F. Neuromuscular fatigue and recovery in elite female soccer: effects of active recovery. *Med Sci Sports Exerc* 2008; 40: 372–380.
  16. Kono H, Chen CJ, Ontiveros F, Rock KL. Uric acid promotes an acute inflammatory response to sterile cell death in mice. *J Clin Invest* 2010; 120: 1939–1949.
  17. Fathallah-Shaykh SA, Cramer MT. Uric acid and the kidney. *Pediatr Nephrol* 2014; 29: 999–1008.
  18. Martillo MA, Nazzari L, Crittenden DB. The crystallization of monosodium urate. *Curr Rheumatol Rep* 2014; 16: 400.
  19. Strowig T, Henao-Mejia J, Elinav E, Flavell R. Inflammasomes in health and disease. *Nature* 2012; 481: 278–286.
  20. Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006; 440: 237–241.
  21. Srinivasula SM, Poyet JL, Razmara M, Datta P, Zhang Z, Alnemri ES. The PYRIN-CARD protein ASC is an activating adaptor for caspase-1. *J Biol Chem* 2002; 277: 21119–21122.
  22. Ren K, Torres R. Role of interleukin-1 $\beta$  during pain and inflammation. *Brain Res Rev* 2009; 60: 57–64.
  23. Li WW, Guo TZ, Liang D, Shi X, Wei T, Kingery WS, Clark JD. The NALP1 inflammasome controls cytokine production and nociception in a rat fracture model of complex regional pain syndrome. *Pain* 2009; 147: 277–286.
  24. Daheshia M, Yao JQ. The interleukin 1 $\beta$  pathway in the pathogenesis of osteoarthritis. *J Rheumatol* 2008; 35: 2306–2312.
  25. Qi J, Ye X, Ren G, Kan F, Zhang Y, Guo M, Zhang Z, Li D. Pharmacological efficacy of anti-IL-1 $\beta$  scFv, Fab and full-length antibodies in treatment of rheumatoid arthritis. *Mol Immunol* 2014; 57: 59–65.
  26. So A, De Smedt T, Revaz S, Tschopp J. A pilot study of IL-1 inhibition by anakinra in acute gout. *Arthritis Res Ther* 2007; 9: R28.
  27. Kawasaki Y, Xu ZZ, Wang X, Park JY, Zhuang ZY, Tan PH, Gao YJ, Roy K, Corfas G, Lo EH, Ji RR. Distinct roles of matrix metalloproteases in the early- and late-phase development of neuropathic pain. *Nat Med* 2008; 14: 331–336.
  28. Ruohonen S, Khademi M, Jagodic M, Taskinen HS, Olsson T, Roytta M. Cytokine responses during chronic denervation. *J Neuroinflammation* 2005; 2: 26.
  29. Hashizume H, DeLeo JA, Colburn RW, Weinstein JN. Spinal glial activation and cytokine expression after lumbar root injury in the rat. *Spine* 2000; 25: 1206–1217.
  30. Guo W, Wang H, Watanabe M, Shimizu K, Zou S, LaGraize SC, Wei F, Dubner R, Ren K. Glial-cytokine-neuronal interactions underlying the mechanisms of persistent pain. *J Neurosci* 2007; 27: 6006–6018.
  31. Busso N, Ea HK. The mechanisms of inflammation in gout and pseudogout (CPP-induced arthritis). *Reumatismo* 2012; 63: 230–237.
  32. Clavijo-Cornejo D, Martínez-Flores K, Silva-Luna K, Martínez-Nava G A, Fernández-Torres J, Zamudio-Cuevas Y, Guadalupe Santamaría-Olmedo M, Granados-Montiel J, Pineda C, López-Reyes A. The overexpression of NALP3 inflammasome in knee osteoarthritis is associated with synovial membrane prolidase and NADPH oxidase 2. *Oxid Med Cell Longev* 2016; 2016: 1472567.
  33. Shinoda M, Ozaki N, Sugiura Y. Involvement of ATP and its receptors on nociception in rat model of masseter muscle pain. *Pain* 2008; 134: 148–157.
  34. Ide S, Minami M, Ishihara K, Uhl GR, Satoh M, Sora I, Ikeda K. Abolished thermal and mechanical antinociception but retained visceral chemical antinociception induced by butorphanol in mu-opioid receptor knockout mice. *Neuropharmacology* 2008; 54: 1182–1188.
  35. Ju TJ, Dan JM, Cho YJ, Park SY. Inhibition of inducible nitric oxide synthase attenuates monosodium urate-induced inflammation in mice. *Korean J Physiol Pharmacol* 2011; 15: 363–369.
  36. Schmidt AP, Bohmer AE, Antunes C, Schallenberger C, Porciuncula LO, Elisabetsky E, Lara DR, Souza DO. Anti-nociceptive properties of the xanthine oxidase inhibitor allopurinol in mice: role of A1 adenosine receptors. *Br J Pharmacol* 2009; 156: 163–172.
  37. Wang S, Li Y, Song X, Wang X, Zhao C, Chen A, Yang P. Febuxostat pretreatment attenuates myocardial ischemia/reperfusion injury via mitochondrial apoptosis. *J Transl Med* 2015; 13: 209.
  38. Wang S, Zhao J, Wang H, Liang Y, Yang N, Huang Y. Blockage of P2X7 attenuates acute lung injury in mice by inhibiting NLRP3 inflammasome. *Int Immunopharmacol* 2015; 27: 38–45.
  39. Xia M, Boini KM, Abais JM, Xu M, Zhang Y, Li PL. Endothelial NLRP3 inflammasome activation and enhanced neointima formation in mice by adipokine visfatin. *Am J Pathol* 2014; 184: 1617–1628.

40. Weisser SB, van Rooijen N, Sly LM. Depletion and reconstitution of macrophages in mice. *J Vis Exp* 2012; 4105.
41. Horai R, Asano M, Sudo K, Kanuka H, Suzuki M, Nishihara M, Takahashi M, Iwakura Y. Production of mice deficient in genes for interleukin (IL)-1alpha, IL-1beta, IL-1alpha/beta, and IL-1 receptor antagonist shows that IL-1beta is crucial in turpentine-induced fever development and glucocorticoid secretion. *J Exp Med* 1998; 187: 1463–1475.
42. Yang G, Yeon SH, Lee HE, Kang HC, Cho YY, Lee HS, Lee JY. Suppression of NLRP3 inflammasome by oral treatment with sulforaphane alleviates acute gouty inflammation. *Rheumatology (Oxford, England)* 2018; 57: 727–736.
43. Lee HE, Yang G, Kim ND, Jeong S, Jung Y, Choi JY, Park HH, Lee JY. Targeting ASC in NLRP3 inflammasome by caffeic acid phenethyl ester: a novel strategy to treat acute gout. *Sci Rep* 2016; 6: 38622.
44. Bullon P, Alcocer-Gomez E, Carrion AM, Marin-Aguilar F, Garrido-Maraver J, Roman-Malo L, Ruiz-Cabello J, Culic O, Ryffel B, Apetoh L, Ghiringhelli F, Battino M, Sanchez-Alcazar JA, Cordero MD. AMPK phosphorylation modulates pain by activation of NLRP3 inflammasome. *Antioxid Redox Signal* 2016; 24: 157–170.
45. Cordero MD, Alcocer-Gómez E, Marín-Aguilar F, Rybkina T, Cotán D, Pérez-Pulido A, Alvarez-Suarez JM, Battino M, Sánchez-Alcazar JA, Carrión AM, Culic O, Navarro-Pando JM, Bullón P. Mutation in cytochrome b gene of mitochondrial DNA in a family with fibromyalgia is associated with NLRP3-inflammasome activation. *J Med Genet* 2016; 53: 113–122.
46. Zhang ZT, Du XM, Ma XJ, Zong Y, Chen JK, Yu CL, Liu YG, Chen YC, Zhao LJ, Gc L. Activation of the NLRP3 inflammasome in lipopolysaccharide-induced mouse fatigue and its relevance to chronic fatigue syndrome. *J Neuroinflammation* 2016; 13: 71.
47. Ding M, Huang T, Zhu R, Gu R, Shi D, Xiao J, Guo M, Li J, Hu J, Liao H. Immunological behavior analysis of muscle cells under IFN-gamma stimulation in vitro and in vivo. *Anat Rec* 2018; 301: 1551–1563.
48. McBride MJ, Foley KP, D'Souza DM, Li YE, Lau TC, Hawke TJ, Schertzer JD. The NLRP3 inflammasome contributes to sarcopenia and lower muscle glycolytic potential in old mice. *Am J Physiol Endocrinol Metab* 2017; 313: E222–e232.
49. Terlizzi M, Colarusso C, Popolo A, Pinto A, Sorrentino R. IL-1alpha and IL-1beta-producing macrophages populate lung tumor lesions in mice. *Oncotarget* 2016; 7: 58181–58192.
50. Li X, Zhang Y, Jin Q, Xia KL, Jiang M, Cui BW, Wu YL, Song SZ, Lian LH, Nan JX. Liver kinase B1/AMP-activated protein kinase-mediated regulation by gentiopicroside ameliorates P2X7 receptor-dependent alcoholic hepatosteatosis. *Br J Pharmacol* 2018; 175: 1451–1470.
51. Gregory NS, Brito RG, Fusaro MC, Sluka KA. ASIC3 is required for development of fatigue-induced hyperalgesia. *Mol Neurobiol* 2016; 53: 1020–1030.
52. Aibibula Z, Ailixiding M, Iwata M, Piao J, Hara Y, Okawa A, Asou Y. Xanthine oxidoreductase activation is implicated in the onset of metabolic arthritis. *Biochem Biophys Res Comm* 2016; 472: 26–32.
53. Wan X, Xu C, Lin Y, Lu C, Li D, Sang J, He H, Liu X, Li Y, Yu C. Uric acid regulates hepatic steatosis and insulin resistance through the NLRP3 inflammasome-dependent mechanism. *J Hepatol* 2016; 64: 925–932.
54. Ives A, Nomura J, Martinon F, Roger T, LeRoy D, Miner JN, Simon G, Busso N, So A. Xanthine oxidoreductase regulates macrophage IL1beta secretion upon NLRP3 inflammasome activation. *Nat Commun* 2015; 6: 6555.
55. Hande KR, Noone RM, Stone WJ. Severe allopurinol toxicity. Description and guidelines for prevention in patients with renal insufficiency. *Am J Med* 1984; 76: 47–56.
56. Beara-Lasic L, Pillinger MH, Goldfarb DS. Advances in the management of gout: critical appraisal of febuxostat in the control of hyperuricemia. *Int J Nephrol Renovasc Dis* 2010; 3: 1–10.
57. Sluyter R. The P2X7 receptor. *Adv Exp Med Biol* 2017; 1051: 17–53.
58. Wu B, Peng L, Xie J, Zou L, Zhu Q, Jiang H, Yi Z, Wang S, Xue Y, Gao Y, Li G, Liu S, Zhang C, Li G, Liang S, Xiong H. The P2X7 receptor in dorsal root ganglia is involved in HIV gp120-associated neuropathic pain. *Brain Res Bull* 2017; 135: 25–32.
59. Zhong X, Zhu F, Qiao J, Zhao K, Zhu S, Zeng L, Chen X, Xu K. The impact of P2X7 receptor antagonist, brilliant blue G on graft-versus-host disease in mice after allogeneic hematopoietic stem cell transplantation. *Cell Immunol* 2016; 310: 71–77.
60. Naithani P, Vashisht N, Khanduja S, Sinha S, Garg S. Brilliant blue G-assisted peeling of the internal limiting membrane in macular hole surgery. *Indian J Ophthalmol* 2011; 59: 158–160.
61. Chen L, Li X, Huang L, Wu Q, Chen L, Wan Q. Chemical stimulation of the intracranial dura activates NALP3 inflammasome in trigeminal ganglia neurons. *Brain Res* 2014; 1566: 1–11.
62. Hussain S, Sangtian S, Anderson SM, Snyder RJ, Marshburn JD, Rice AB, Bonner JC, Garantzotis S. Inflammasome activation in airway epithelial cells after multi-walled carbon nanotube exposure mediates a profibrotic response in lung fibroblasts. *Part Fibre Toxicol* 2014; 11: 28.
63. Kim MJ, Yoo JY. Active caspase-1-mediated secretion of retinoic acid inducible gene-I. *J Immunol* 2008; 181: 7324–7331.
64. Zaslonka Z, Palsson-McDermott EM, Menon D, Haneklaus M, Flis E, Prendeville H, Corcoran SE, Peters-Golden M, O'Neill L. The induction of pro-IL-1beta by lipopolysaccharide requires endogenous prostaglandin E2 production. *J Immunol (Baltimore)* 2017; 198: 3558–3564.
65. Yazdi AS, Drexler SK. Regulation of interleukin 1alpha secretion by inflammasomes. *Ann Rheum Dis* 2013; 72: ii96–ii99.
66. Groß O, Yazdi AS, Thomas CJ, Masin M, Heinz LX, Guarda G, Quadroni M, Drexler SK, Tschopp J. Inflammasome activators induce interleukin-1alpha secretion via distinct pathways with differential requirement for

- the protease function of caspase-1. *Immunity* 2012; 36: 388–400.
67. Oguri S, Motegi K, Iwakura Y, Endo Y. Primary role of interleukin-1 alpha and interleukin-1 beta in lipopolysaccharide-induced hypoglycemia in mice. *Clin Diagn Lab Immunol* 2002; 9: 1307–1312.
68. Vonk AG, Netea MG, van Krieken JH, Iwakura Y, van der Meer JW, Kullberg BJ. Endogenous interleukin (IL)-1 alpha and IL-1 beta are crucial for host defense against disseminated candidiasis. *J Infect Dis* 2006; 193: 1419–1426.
69. Sjogaard G, Sogaard K. Muscle activity pattern dependent pain development and alleviation. *J Electromyogr Kinesiol* 2014; 24: 789–794.